Implementation of a Social Needs Screening Tool Across Multiple Settings for Children with Asthma

Shayla Stringfield, BS, CHES^{1,2}; Taylor Brewer, BS²; Rachel Margolis, PhD, LICSW^{1,3}; Amanda Jepson, BS⁵; Kavita Parikh, MD, MSHS⁵; Shilpa J. Patel, MD, MPH^{1,4} ¹IMPACT DC Asthma Clinic; ²George Washington University School of Medicine and Health Sciences; ³Center for Translational Research, Children's Research Institute; ⁴Division of Emergency Medicine, ⁵Division of Hospital Medicine Children's National Health System; Washington, DC

BACKGROUND


- Social determinants of health contribute to higher rates of asthma morbidity and health disparities in urban populations.
- If feasible, a standardized approach to address social needs across a health system could prove efficient and improve outcomes.


OBJECTIVES

REFERENCES

- To implement standardized screening for unmet social needs among families with children with asthma
- To **assess** feasibility (proportion screened) of standardized screening across different settings
- To **describe** unmet social needs, indicated by assistance requested RESULTS on screening tool (Figure 1)

METHODS





Care Unit (PICU)*

(SAC)

N=299


•Stempel H, Federico MJ, Szefler SJ. Applying a biopsychosocial model to inner city asthma: Recent approaches to address pediatric asthma health disparities. Pediatric Respiratory Reviews. Published online January 1, 2019. doi:10.1016/j.prrv.2019.07.001

•Teach S.J., Crain EF, Quint D.M. Hylan ML, Joseph J.G. Improved Asthma Outcomes in a High-Morbidity Pediatric Population: Results of an Emergency Department-Based Randomized Clinical Trial. Arch Pediatr Adolesc Med. 2006; 160(5):535. doi:10.1001/archpedi.160.5.535 Margolis, R. H. F., Patel, S. J., Krueger, J., Brewer, T., Williams, A., Stringfield, S., Teach, S. J., & Parikh, K. (2023). Association between social needs and asthma control among children evaluated at a single-center high-risk asthma clinic. The journal of allergy and clinical immunology. In practice, \$2213-2198(23)00286-6. Advance online publication. https://doi.org/10.1016/j.jaip.2023.03.0043.

RESULTS

 The PICU continued screening following the pilot while the SAC and the ED did not. Across the three locations and asthma clinic, 42% (165/390) of families requested help with ≥1 social need.

Figure 3: Distribution of Unmet Social Needs



CONCLUSION

Screening was feasible in some locations, more successful in higher risk patients, and overall revealed a high burden of social needs among families of children with asthma across our institution.




Figure 1: Social needs screening tool







